

REPORT

DISTRIBUTION OF NEOTROPICAL OTTER, *Lontra longicaudis* (OLFERS, 1818) (MUSTELIDAE) IN COASTAL ISLANDS OF SANTA CATARINA, SOUTHERN BRAZIL

Oldemar CARVALHO-JUNIOR^{1*}, Alexandre FILLIPINI², Carlos SALVADOR³

^{1*}Instituto Ekko Brasil, IEB, Brazil e-mail: atendimento@ekkobrasil.org.br

²IBAMA, Florianópolis, SC, Brazil

³Cooperativa Caipora, Florianópolis, SC, Brazil



(received 11th April 2012, accepted 13th July 2012)

Abstract: This study presents the distribution of neotropical otters (*Lontra longicaudis*) in coastal islands of Santa Catarina State. A total of 23 islands were surveyed, through 46 field trips. Otter's signs were found in 9 islands, representing 39% of the total. It is believed that the *Lontra longicaudis* use these islands as a support for its movement from one place to another. The results could be indicating the existence of a coastal ecological corridor used by the neotropical otter.

Keywords: Ecology of otter, occurrence, coastal habitats, ecological corridor

INTRODUCTION

Distribution of *Lontra longicaudis* in marine habitats along the coast of Brazil is poorly known. The neotropical otter in Brazil is often registered in rivers, estuaries, mangroves, lakes and lagoons (Carvalho-Junior, 1990; Helder and Andrade, 1997; Pardini, 1998; Quadros and Monteiro-Filho, 2001; Alarcon and Simões-Lopes, 2004; Quintela et al., 2008; Carvalho-Junior et al., 2010a,b). However, citations of neotropical otter on coastal islands are rare.

Santa Catarina state has at least 130 islands, Santa Catarina Island (Florianópolis) being the largest one. The island of Santa Catarina has an average length of 54 km and width average of 18 km. It has several bays, promontories, islands, bays and lagoons. The island is located parallel to the mainland and separated by a narrow channel.

Some islands are distant from the mainland such as Badejo Island (21.2 km), Xavier Island (17.8 km), Mata-Fome Island (16.4 km), Campeche Island (16.0 km) and Arvoredo Island (12.62 km). If taking into account the presence of Santa Catarina Island, some distances can be reduced. Campeche Island, for example, is 16 km away from the mainland, but only 1.4 km distant from Santa Catarina Island. Arvoredo

Island, on the other hand, is located north of Santa Catarina Island and its distance from mainland and from Santa Catarina Island is similar, 10.6 km.



Figure 1. Landsat satellite photo of Santa Catarina Island (Source: <https://zulu.ssc.nasa.gov/m>). It can be observed two main lagoons, Peri Lake at the South and Conceição Lagoon at the North.

Presence of *Lontra longicaudis* on islands, close to the mainland of Santa Catarina state (less than 1 km), has been reported before. Anhatomirim Island (Alarcon and Simões-Lopes, 2004) and Porto Belo Island (João da Cunha Island) (Carvalho-Junior, 2007) are known for the presence of the species throughout the year. However, a comprehensive study of the distribution of the species on coastal islands was never carried out before.

Carvalho-Junior et al. (2004) studied the distribution and characteristics of environments used by the neotropical otter in the coastal region of Santa Catarina State, Brazil. What is important to point out is the indented coastline of the state. It is replete of small bays and promontories that in turn can result in different distances between mainland and islands, depending on the angle the observer takes.

The coastal area of Santa Catarina is also rich in lakes, lagoons and river mouths. The largest number of lagoons in mainland is located between Arvoredo Island (27°17'98" S/48°21'53" W) and Lobos Island (28°26'48" S/48°42'33" W). This

area has a number of lakes that includes (from south to north), the Camacho Lagoon (6.32 km²), Santo Antonio Lagoon (33.85 km²), Imaruí Lagoon (86.32 km²), Mirim Lagoon (63.77 km²), Ibiraquera Lagoon (8.65 km²) and Garopaba Lagoon (5.15 km²), along with several rivers flowing into the sea.

In Santa Catarina Island the most important water bodies are the Peri Lake (5.1 km²) and the Conceição Lagoon (19.71 km²) (Figure 1). To the North of the Santa Catarina Island is more common to find mouths of rivers like Tijucas River, Itajai-Açú, Itapocu River, and Babitonga Bay in the extreme north of the state. The presence of promontories is also significant such as Penha, Porto Belo and Governador Celso Ramos Promontories (Figure 2).




Figure 2. CBERS Satellite Image of the Coastline of Santa Catarina State. (a) North, (b) Central, (c) South. (Courtesy: INPE - National Institute of Spatial Research).

Distribution of otters on coastal islands, further away of the continent, such as Campeche Island and Irmã-do-Meio Island, has been reported only recently (Carvalho-Junior, 2007), suggesting that the presence of the *Lontra longicaudis* in marine environments could be more intense than expected. This work represents a register of the presence of *Lontra longicaudis* in coastal islands of Santa Catarina State located between 26°42'33" S and 28° 32' 21" S (Figure 3).



Figure 3. CBERS Satellite Image of the Coastline of Santa Catarina State. (a) North, (b) Central, (c) South. (Courtesy: INPE - National Institute of Spatial Research). The study area included 23 islands from 26°42'33" S to 28°32'21" S. (1. Itacolomis Island, 2. Feia Island, 3. Deserta Island, 4. Arvoredo Island, 5. Ratonés-Grande Island, 6. Guarás-Grande Island, 7. Guarás-Pequeno Island, 8. Mata-Fome Island, 9. Badejo Island, 10. Xavier Island, 11. Campeche Island, 12. Irmã-do-Meio Island, 13. Moleques-do-Sul Island, 14. Cardos Island, 15. Corais Island, 16. Siriú Island, 17. Batuta Island, 18. Santana-de-Dentro Island, 19. Santana-de-Fora Island, 20. Araras Island, 21. Tacami Island, 22. Lobos Island, 23. Ilhota).

MATERIALS AND METHODS

Each island was inspected on foot, from 6:00 AM to 06:00 PM, searching for signs of otters. Indirect signs of the presence of otters were considered to be footprints, feces, scent marks (urine and mucus) and the presence of active shelters. The islands were visited twice, fall/winter and spring/summer 2007/2008.

The organization and planning of field trips included preliminary data collection, such as maps, aerial photos, satellite images and selection of the islands to be visited. The study area chosen was the one with the islands scattered in the central coast of Santa Catarina. The 23 islands and their locations, selected for research can be seen at (Table 1).

The fieldwork on each island was finalized after the entire surface had been explored. The time spent on each island varied depending on the size and difficulty of movement due to density of vegetation and topography.

Small islands, flat and smooth, as Guarás-Pequeno, Guarás-Grande, Cardoso, Itacolomis, Santanas and Ilhota Islands, were examined from 30 minutes to 3 hours. Larger islands such as Arvoredo, Irmã-de-Dentro, Moleques-do-Sul and Corais imposed a full day's work, resulting in 1 to 3 days of fieldwork.

A total of 23 field trips were conducted in the spring/summer, totaling 127 hours worked, and 21 field trips in the fall/winter, with 105 hours worked. The total effort was 44 visits, corresponding to 232 hours of research.

Table 1. Surveyed Islands and coordinates.

Island	Coordinates
1. Itacolomis	26°42'35" S/48°37'05" W
2. Feia	26°44'41" S/48°38'11" W
3. Arvoredo	27°17'98" S/48°21'53" W
4. Deserta	27°16'17" S/48°19'50" W
5. Campeche	27°41'49" S/48°27'52" W
6. Xavier	27°33'09" S/48°35'05" W
7. Mata-Fome	27°25'29" S/48°22'00" W
8. Badejo	27°26'33" S/48°20'49" W
9. Ratonés-Grande	27°42'30" S/48°35'44" W
10. Guarás-Grande	27°33'09" S/48°35'05" W
11. Guarás-Pequeno	27°32'58" S/48°33'09" W
12. Irmã-do-Meio	27°50'19" S/48°31'33" W
13. Moleques-do-Sul	27°50'45" S/48°25'53" W

14. Cardos	27°48'52" S/48°34'51" W
15. Siriú	27°59'15" S/48°37'04" W
16. Corais	27°56'13" S/48°32'32" W
17. Tacami	28°21'06" S/48°36'04" W
18. Santana-de-Dentro	28°14'28" S/48°39'06" W
19. Santana-de-Fora	28°14'53" S/48°39'58" W
20. Batuta	28°09'10" S/48°38'32" W
21. Araras	28°19'17" S/48°38'55" W
22. Lobos	28°26'48" S/48°42'33" W
23. Ilhota	28°32'21" S/48°45'35" W

RESULTS

According to Filippini (2009), in the 23 islands, 14 different habitats were classified: supralittoral rocky, supralittoral sandy, mesolittoral rocky, mesolittoral sandy, herbaceous vegetation, shrub vegetation, arboreal vegetation, disturbed vegetation, wetland, ground, cave, lagoon, creek and edification.

All 23 islands surveyed showed the habitat supralittoral rocky, dominated by rock, absence of vegetation, subjected to insulation, rain, wind and marine spray. All islands have rocky supralittoral zone. *Lontra longicaudis* signs were found in this habitat, together with *Zonotrichia capensis*, *Pitangus sulphuratus*, *Chatartes sp.*, *Tupinambis sp.*, *Sula leucogaster*, *Phalacrocorax brasilianus*, *Otaria flavescens*, *Arctocephalus australis*, domestic goat, *Larus dominicanus*, and *Thalasseus sp.* In total, 40 species of vertebrates were found in the supralittoral rocky.

The supralittoral sandy is the habitat formed between the end of the vegetation and the high tide. This habitat was found in five islands: Ratonés-Grande, Guarás-Grande, Guarás-Pequeno, Campeche and Corais. Signs of neotropical otter were not found in this region. Other vertebrates observed in this habitat consist mainly of *Larus dominicanus*, *Coragyps atratus*, and domestic goats.

The mesolittoral rocky zone represents the rocky surface covered by marine benthos, placed between the maximum high tide and the low tide. All 23 surveyed islands have rocky mesolittoral zone. It was found 20 vertebrate inhabiting this zone. The typical species is the *Haematopus palliatus*, adapted to feed on molluscs in the rocks. Other vertebrates are the *Larus dominicanus*, *Zonotrichia capensis*, *Troglodytes musculus*, *Pitangus sulphuratus*, *Ceryle torquata*, *Charadrius semipalmatus*, three species of Scolopacidae and five species of the Ardeidae family. If the otter was observed in the supralittoral zone we can also deduce that this area is used by the species.

The mesolittoral sandy is the habitat formed by the intertidal action in the sandy beaches. The islands that presented mesolittoral sandy were: Ratonés-Grande, Guarás-Grande, Guarás-Pequeno, Campeche and Corais. It was observed five species in this region: *Larus dominicanus*, *C. atratu*, *Haematopus palliatus*, *Polyborus plancus* and *Zonotrichia capensis*. As for the supralittoral sandy, signs of neotropical otter were not found in this region.

The herbaceous vegetation habitat is formed predominantly by herbaceous vegetation with up to a meter height. It is present in 14 islands and appears along the cliffs, flat areas and wetlands, overlying sandy deposits. The most common plants are *Ipomoea*, *Panicum*, *Paspalum*, *Spartina*, *Remirea*, *Lantana*, *Tibouchina*, *Vitex*, *Aechmea*, *Vriesea*, *Cereus*, *Opuntia*, and *Polypodium*. It was observed 34 species of vertebrates in this habitat, including *Larus dominicanus* and *Sula leucogaster*. It is also common to see forest birds, such as *G. aequinoctialis*, *B. culicivorus*, *Troglodytes musculus*, *Zonotrichia capensis*, *Pitangus sulphuratus*, *Coragyps atratus*, *Speotyto cunicunaria*, *Milvago chimango*, *Milvago chimachima*, *Polyborus plancus*, *Rupornis magnirostris* and *Syrigma sibilatrix*. Other species are lizards, snakes, rabbits and rodents Cavidae.

The shrub vegetation habitat consists of shrubby plants with 1 to 5 meters tall, observed in 15 islands. As for herbaceous vegetation, it is found overlying sandy deposits along the coasts. The main species of vegetation are *Schinus terenetifolius*, *Ocotea sp.*, *Butia capitata*, *Eugenia sp.*, *Myrcia sp.*, *Vitex sp.*, *Campomanesia sp.*, *Tibouchina sp.*, *Guapira opposita*, *Myrsine sp.*, *Psidium sp.*, *Tabebuia sp.*, *Cereus* and *Opuntia*. It was observed 42 species in this habitat, consisted mainly of birds adapted to a more diverse plant condition. *Tyrannus melancholicus* and the *Guira guira* are frequent in this habitat. Presence of birds adapted to shade and humidity of trees were also registered, such as *Turdus sp.* (Muscicapidae), *Elaenia sp.* (Tyrannidae), *Vireo chivi* (Vireonidae), *Ranphocelus bresileus*, *Thraupis sayaca*, and *Coereba flaveola* (Emberizidae).

Trees with height greater than 0.5 m form the arboreal vegetation habitat. The main species are *Clusia sp.*, *Syagrus romanzofiana*, *Alchornea sp.*, *Ficus sp.*, *Coussapoa sp.*, *Inga sp.*, *Nectandra sp.*, *Ocotea sp.*, *Psidium sp.*, *Ilex sp.*, *Cecropia sp.*, *Myrsine sp.*, *Eugenia sp.*, *Guapira opposita*, *Gomidesia sp.*, *Myrcia sp.*, *Citharexylum mirianthum*, *Cupania sp.*, *Tabebuia sp.*, *Hibiscus sp.* and *Huberia sp.* The largest number of vertebrates, with 48 species, inhabits this habitat. The typical species are those adapted to shadows and humidity of the forest, such as Chiroptera, Muscicapidae, Tyrannidae, Colubridae, Hylideos and *Didelphis sp.*

The disturbed vegetation habitat is formed by exotic vegetation. Often this habitat is close to buildings. It was found in 9 islands and it is inhabited by 30 vertebrates, constituted basically by native and domestic introduced animals as birds (*Troglodytes musculus*, *Zonotrichia capensis* and *Pitangus sulfuratus*), lizards (*Tabebuia sp.*), rooster, ducks, goose, chickens, cats and dogs.

The wetland habitat is formed in depressions due to the accumulation of rainwater or groundwater, accompanied by vegetation adapted to flooding. The most common species are the grasses *Paspalum sp.*, *Spartina sp.*, *Scirpus sp.*, *Cyperus scleria* and *Typha sp.* It was found 4 wetlands in 3 islands, the Mata-Fome, Xavier and Corais. Three of these habitats were in the high parts of the islands and a fourth was located near sea level. It was found only amphibious using this habitat.

The ground habitat consists of the soil or substrate formed on the basement rock, usually covered by vegetation. It was found 3 species using this habitat, a snake *Amphisbaena sp.*, an owl *Speotyto cunicunaria* and a rabbit *Oryctolagus cuniculus*.

The cave habitat represent natural rocky cavities, accessible to man, with variable dimensions, with only temporary short time periods of light, a higher thermal stability inside than the outside and a relative humidity that tends to saturation. Four

caves were found in 3 islands. Arvoredo and Batuta Islands with one cave each, and Feia Island with 2 caves. These caves are located between 2 and 3 meters above sea level, at the interface between the supralittoral rocky and native vegetation. Neotropical otter signs were found in 2 caves, one at Batuta Island and another at Arvoredo Island.

The Batuta Island cave is opened to the North, 3 meters large, 5 meters long, 1 meter high and at 3 meters from sea level. It consists of a narrow corridor with walls and floor of granite. The end is covered with rocks and sediments from which runs a trickle of water drainage. The Arvoredo Island cave is opened to the West, 1 meter large, 5 meters long, 1 meter high and at 1.5 meters from sea level.

No signs of otters were found at the 2 caves of Feia Island. The first is opened to the South, 8 meters large, 60 meters long, 3 meters high and at 2.5 meters from sea level. The second is opened to the North, 6 meters large, 30 meters long, 4 meters high and at 2 meters from sea level.

The lagoon habitat is formed by a calm and shallow water body that maintains a restricted communication with the sea, with a salinity that can vary from almost fresh to hypersaline. This habitat is found only in the Siriú Island. This is a small oval lagoon, 36 meters large, 10 meters long and 1.50 meters deep, located in the center of the Island and surrounded by rocks. The water of this lagoon is formed by mixing of rainwater with seawater that penetrates through a gap of 31 m located on the floor at the east side of the Island. It was found 3 species of crabs in the lagoon, *Chasmagnathus granulata* and *Callinectes sp.* Only 2 species of vertebrates was found in the lagoon, *Larus dominicanus* and *Lontra longicaudis*.

The stream habitat is a small perennial water flow. Two streams were found at Arvoredo Island. One is approximately 550 m long and the other is approximately 1100 m long. Both are perennial and completely hidden in the Atlantic Forest that covers the island. The waters of the two streams are limpid, flowing through rocks, gravel and fallen logs. There were no fish but the crustacean *Macrobrachium sp* is common. This crustacean is one of the favorites in the diet of neotropical otter (Carvalho-Junior, 2010). Signs of *Lontra longicaudis* were found at the longer creek.

The edification habitat is the one constructed by man. It consists of houses, cottages, restaurants, fortifications, sheds, lighthouses, piers, water tanks, septic tanks, solar panels, stairs, walls, bathrooms, toilets, walkways or some kind of ruin. Among the islands surveyed, 11 exhibited some type of building. The island containing the smallest constructed area is the Mata-Fome, a shack of 6 m². The island with the largest constructed area is the Campeche, with 2 restaurants, 6 houses, a lodge, 2 barbecue grills, a bathroom, 2 power houses, a gas tank, 4 water tanks and 3 walkways, summing up all together 1700 m². The Arvoredo Island presents a lighthouse, 4 homes, a lodge, a powerhouse, 9 solar panels, a fuel shed, a warehouse and 13 water tanks. *Lontra longicaudis* was found at Arvoredo and Campeche Islands, but not at the edification areas.

The Islands and the neotropical otter

Itacolomis Islands are two granitic islands separated by a 30 meters channel. They are 6.2 km away from mainland, approximately 25 and 40 meters height, 40 and 65 meters long, and an area of 0.52 ha all together. There is no edification in these islands. The total perimeter is 393 meters; maximum width is 26 and 52 meters each,

with steep topography. The islets are used for research, camping and fishery, but not by the neotropical otter.

These islands are located further north in relation to the other surveyed islands. The surface of the islands is 62.63% rocky. In the upper areas of the southern promontory it is observed herbaceous vegetation (31.33%), consisting of grasses and clumps of cactus, *Opuntia sp.*, *Guapira opposita* and *Spinacia oleracea*. The shrub appears only in the northern promontory, formed by groups of *Opuntia sp.*, covering 6.02% of the surface.

Feia Island is situated 4 km southeast of Itacolomis and at 2 km from the coast. It is mountainous, has no buildings, the area is almost 11 ha and is 80 m high. Its perimeter of 1194 meters is very irregular and difficult to walk. The island is 500 m long and a maximum width of 375 m, with an area of 10.95 ha. It has 70% of its area covered by vegetation. It is used for research, camping, visitation and fishing. Most of the coastal perimeter (6.32% of mesolittoral and 14.43% of supralittoral) presents several vertical cracks. Two of these cracks were considered caves, inhabited by *Streptoprogne zonaris*. *Lontra longicaudis* was not registered on this island.

Deserta Island has 15.30 ha and is one of the most distant from the coast (15.7 km) and at 13,7 km from Santa Catarina Island. The east coast is steep and abrupt with 80 m height and a perimeter of nearly 4 km. The island has no buildings. Signs of neotropical otter were found at the west side of the island. The island is covered by 22.22% of clumps of *Guapira opposita*, 27% by herbaceous vegetation of grasses and bromeliads. The remaining area is 15.13% of mesolittoral rocky and 35.63% of supralittoral rocky.

Arvoredo Island is one of the islands where signs of *Lontra longicaudis* were found. It is mountainous, 300 m high, a length of 3.5 km, 1740 m width and a perimeter of 10.5 km. It has an area of 324 ha and is one of the largest islands surveyed. This island is located at 12.6 km from mainland and at 10.7 km from Santa Catarina Island. It presents a variety of environments such as rocky coast, small streams of water, Atlantic Forest, dunes, and buildings, such as lighthouse, houses and sheds.

This island exhibits two unique habitats among all islands surveyed: the Atlantic Forest, which covers 80.76% of the surface of the island, and two perennial streams (0.02%). *Lontra longicaudis* uses both habitats, at the north of the Island. No fishes were found in these 2 streams. During the survey 36 species of vertebrate were found, including *Didelphis aurita*, *Nasua nasua*, *Tupinambis merianae*, *Scinax perereca*, a variety of seabirds (e.g. *Sula leucogaster* and *Fregata magnificens*), birds such as *Egretta thula*, *Butorides striatus*, hawks and forest birds, and domestic species such as dogs and *Capra aegagrus hircus*.

Mata-Fome Island is a small island with 3.49 ha, 40 m high and a perimeter of 1225 m. Is relatively close to Santa Catarina Island (690 m) but 16.4 km away from the mainland. It has a small hut used by fishermen. In the central and highest portion of the island, there is a wetland with *Typha dominigensis*. No amphibians or other vertebrates were found in this habitat. Herbaceous grasses and bromeliads compose the vegetation; however the supralittoral rocky habitat is predominant (48%).

Badejo Island is located 2.4 km southeast of the Mata-Fome Island, 1 km north of the of Santa Catarina Island and 21.2 kilometers from the continent. The island is relatively small (2.84 ha), with steep topography and low altitude (35 m). It is 333 m long, with a perimeter of 1168 m and maximum width of 155 m. It is used for research, camping, fishing and visitation. The island is rocky with meso and supralittoral representing 66.89% of the surface. Areas with herbaceous vegetation and shrubs are 33.14%. Vertebrates found in the island were birds (*Sula leucogaster*, *Coragyps atratus*, *Milvago chimachima*, *Milvago chimango*, *Haematopus palliatus*, *Larus dominicanus*, *Sterna hirundinacea*, *Pitangus sulphuratus*, *Troglodytes aedon*, *Zonotrichia capensis*) and one species of lizard *Hemidactylus mabouya*.

Ratones Grande Island is located at 4.6 km from mainland and 2.3 km from Santa Catarina Island. It presents an area of 21.22 ha, 1 km long, 280 m width and 44 m height. The total perimeter is 2726 meters, mainly rocky. The island is covered by Atlantic Forest and the main building is a historic building, the Fortress of Santo Antonio, built in 1740. It was found 33 vertebrate, including 53 species of birds, 5 mammals, 5 reptiles and 1 amphibian. Among the mammals were the opossum *D. aurita*, a bat *M. nigricans*, and the *Lontra longicaudis*.

The **Guarás-Pequeno Island** was the smallest island with 0.25 ha, 1.7 m height, length of 45 m and a perimeter of 512 m. It is situated within the North Bay, 2.4 km from Santa Catarina Island and 4.0 km away of the continent. This island is used mainly for camping and fishing. The only edification is a barbecue place. The perimeter of the island is rocky with a short sandy beach of 5 meters length. The centre of the island is covered by shrub. Mammals were not found on the island.

The **Guarás-Grande Island** is also located in the North Bay, 2.4 km north of Santa Catarina Island and 3.7 kilometers from mainland. It has 1.1 ha, 5 m height and 171 m long. The west side of the island has small beaches intercalated by rocks. The east side is mainly rocky with some parts of sand. Edifications in the island include 2 warehouses, a house, a kiosk, solar panel, 2 water tanks, 1 chapel and a barbecue. Most of the original vegetation was replaced by exotic vegetation where chickens and dogs walk freely. Neotropical otter was not found in the island. The island is used for military training.

The **Xavier Island** has an area of 8.71 ha and height of 50 m. The topography is steep and the length of 607 m is almost three times its maximum width (219 m). It is located 3.6 km east of Santa Catarina Island. This island is used mainly for research, camping, visitation and fishing, but there are no edifications. The coast of the island is all rocky with no beaches or springs of fresh water. The northern tip is only slightly higher than the rest of the island, allowing the formation of relatively flat areas, including the largest wetland of all the islands surveyed. The shrub vegetation covers 20.37% of the surface while herbaceous covers 48.82%. The terrestrial vertebrate fauna recorded is represented by 17 species, 14 birds, 2 mammals (*Oryctolagus cuniculus* and *Artocephalus australis*) and an exotic reptile (*Hemidactylus mabouya*).

The **Campeche Island** is mountainous, with 53.56 hectares, a perimeter of 5.5 km, 82 meters height, length of 1620 meters and maximum width of 560 meters. It presents rocky shores and a sandy beach of 500 meters long. This island is located at 1415 m east of Santa Catarina Island and 16 kilometers from mainland. It is covered by Atlantic Forest. The buildings (7 houses, bathrooms, 2 restaurants, 4 water tanks, barbecue places) cover approximately 1700 m², 0.31% of the surface of the island. The trees represent 58.51% of the area, followed by shrubs with 11.89%. This island is used for research, fishing, visitation, environmental education, and commerce. In the rocky supralittoral zone 3 latrines of neotropical otter were found. Different species of birds and mammals such as *Didelphis sp*, *Nasua nasua*, *Dasyprocta azarae*, *Mus musculus* and *Arctocephalus sp*. are also part of the fauna in the island.

Cardoso Island is located in the South Bay, 1.5 km away from Santa Catarina Island and 1.23 km from the continent. The relief is undulated with approximately 8 m high, 145 m in length, a perimeter of 358 meters, maximum width of 80 meters and an area of 0.78 ha. It is used for signaling boating, camping, fishing and research. The meso and supralittoral rocky represent 43.22% of the area of the island while the herbaceous vegetation covers 56.76% of the surface. The terrestrial vertebrate fauna recorded on the island consists of 13 species, 11 birds, a mammal and an exotic reptile. Signs of *Lontra longicaudis* were not found on this island.

Irmã-do-Meio Island presents an area of 55 ha and a height of 103 m. The Atlantic Forest covers 67.60% of its surface. The remaining is 14.75% of shrub distributed on the exposed side (east) and 0.26% of disturbed vegetation located on the north side of the island. The island has no permanent water and no edifications. It is used for fishing, camping and visitation. It was recorded 33 species of vertebrates, 27 birds, 3 mammals (*Didelphis aurita*, a bat and the neotropical otter), two reptiles (*Tupinambis merianae*, *Hemidactylus mabouya*) and a frog (*Leptodactylus cf. gracilis*). In the nearly 6 km of coastline, the rocky supralittoral, it was verified the presence of otters, two latrines and a cave.

Moleques-do-Sul Island has no beach and no cave. The island has an area of 10.6 ha with maximum width of 275 m. The perimeter is 2464 m, length of 705 m and 115 m high. It is distant 14000 meters from mainland and 8.237 meters from Santa Catarina Island. The island has no freshwater. The rocky area (supra and mesolittoral) covers 32.10% of the surface of the island. It is the only one of the 23 islands surveyed that has all three habitats of native vegetation (trees, shrubs and herbaceous). The vertebrate fauna includes 29 species, 26 birds, 1 mammal (*Cavia intermedia*) and 1 reptile (*Amphisbaena sp.*). The island is used for fishing, camping, research and collecting mussels. The neotropical otter was not found on the island.

Coral Island is 4.6 km distant from mainland. It has an area of 31.28 ha, length of 1130 m, maximum width of 406 m, height of 65 m, and a perimeter of 3315 m. The terrain is mountainous. It is used for fishing, collecting mussels, visitation, catching birds, camping and nautical signage. The buildings include a lighthouse, a chapel and a staircase. Shrub represents 49.71% of the area and arboreal vegetation 23.60%. There are two wetlands. It was found 35 species, 3 mammals (*Didelphis aurita*,

Arctocephalus australis, *Capra aegagrus hircus*), 27 birds, 3 reptiles and 2 amphibians. Neotropical otter was not found on the island.

Siriu Island is a small island with less than 2 ha, 12 m height, 220 m long and a perimeter of 744 m. The distance from mainland is 1167 m. The lagoon is a small body of water in the central part of the island maintained by rain and seawater. The area is used for fishing, camping, collecting mussels, visitation and research. The herbaceous vegetation accounts for 19.19% of the area. The rocky surface covers 79.10% of the island in which 44.33% is supralittoral and 34.77% is mesolittoral. The island is used by the neotropical otter. Otter feces were found in the rocky supralittoral zone. The vertebrate fauna of the island is 14 species, 11 birds, one mammal (the neotropical otter), a reptile and an amphibian.

The **Batuta Island** is situated at 350 m from the mainland, in front of a lagoon (Ibiraquera Lagoon). It is 40 m height with an area of 8.17 ha, length of 343 m, maximum width of 307 and a perimeter of 1210 m. The island is used for fishing, collecting mussels and visitation. The surface is 64.51% shrubs, 5.35% herbaceous and 0.21% disturbed vegetation. There is a cave on the north shore used by the neotropical otter. It was recorded 29 species, 23 birds, 4 mammals, an amphibian and a reptile.

Santana-de-Dentro Island is very close to the continent (82 m). It has 3.72 ha, a perimeter of almost 1 km, length of 270 m, maximum width of 172 m and 12 m height. Southeast of the island there is a natural salt-water pool with 7 m diameter and 3 m depth. The island is used for fishing, collecting mussels and visitation. There is no edification in the area. The herbaceous vegetation covers 53.57% of the island. The other half is rocky (15.83% supralittoral and 30.58% mesolittoral). It was found 12 species, 11 birds and a mammal, the neotropical otter. Signs of the otter were found along the rocky shore.

Santana-de-Fora Island has 4.59 ha and is located 540 m from the mainland. It is 25 m height, with a perimeter of 1155 m, 277 m of length and width of 197 m. The island is used for fishing, visiting and collecting mussels. There is only one house in the island. The rocky supralittoral represents 28.84% of the area and herbaceous vegetation covers 54.24% of the island. The supralittoral rocky is also used by the neotropical otter. Besides the otter it was registered 14 species of exotic birds and a reptile.

Araras Island is an island with nearly 10 ha, 779 m in length, maximum width of 155 m, maximum height of 44 m and a perimeter of 2512 m. This island is used for fishing, collecting mussels, research and visitation. There is a lighthouse and a cottage. The vegetation is predominantly shrub (27.52%) and herbaceous (13.94%). The rocky supralittoral occupies 40.33% of the island, followed by rocky mesolittoral with 15.63%. It was recorded 19 species, 14 birds, 4 mammals and one reptile. Signs of neotropical otter were not found.

Tacami Island is one of the smaller islands (0.73 ha and 377 m perimeter), 9.6 km

away from the continent. It has a length of 87 m, maximum width of 53 m and 15 m height. It is used for fishing. The herbaceous vegetation occupies only 13.78% of the surface while the rocky area represents 86.21% of the island. Only five species of birds have been recorded. Signs of neotropical otter were not found on the island.

Lobos Island has 8.7 ha, height of 20 m, 400 m in length and maximum width of 284 m. It has a perimeter of 2 km and is 3.7 km away from mainland. It is used for fishing, collecting mussels and camping. The island has a lighthouse for boating signage. There are two dominant habitats, the rocky supralittoral (41.98%) and vegetation (40.79%). It was registered 12 species, 10 birds, a mammal and an exotic reptile. Signs of neotropical otter were not found in this island.

Finally, the **Ilhota Island** has an area of 2.38 ha and it is 15 m high. Has a length of 233 m with a maximum width of 147 m and perimeter of 947 m. It is located at 156 m from the continent. It is used for fishing, collecting mussels, camping and research. The supralittoral rocky represents 44.13% of the total area while the mesolittoral rocky is 30.51%. The vegetation is predominantly shrub (22.62%) and herbaceous (2.72%). It was registered 21 species, 2 mammals and 18 species of birds. Signs of neotropical otter were not found on this island.

Therefore, from the total number of islands visited, 9 (39%) showed signs of otters: Arvoredo, Campeche, Deserta, Ratonés-Grande, Irmã-do-Meio, Siriú, Santana-de-Dentro, Santana-de-Fora e Batuta (Figure 4). Table 2 shows the main habitats in the islands where signs of otters were found.




Figure 4. Islands where presence of the neotropical otter was registered.

Table 2. Islands where the neotropical otters were present and main characteristics.

Island	Characteristics
Arvoredo	Atlantic Forest, fresh-water, caves
Campeche	Atlantic Forest, fresh-water, caves
Deserta	Shrub and herbaceous vegetation
Ratones-Grande	Atlantic Forest
Irmã-do-Meio	Atlantic Forest
Siriú	Herbaceous vegetation

Santana-de-Dentro	Herbaceous vegetation
Santana-de-Fora	Herbaceous vegetation
Batuta	Shrub and herbaceous vegetation, cave

DISCUSSION

Altogether, 98 species of vertebrates were registered on the 22 islands surveyed: 75 birds (76.53%), 14 mammals (14.28%), 6 reptiles (6.1%) and 3 amphibians (3%). The neotropical otter appeared in 9 islands. Places where otters were observed were mainly related to the interface between sea and land.

The presence of otters was related with the rocky supralittoral zone associated with Atlantic Forest, caves and fresh-water (Arvoredo and Campeche Islands), with Atlantic Forest (Ratones-Grande and Irmã-do-Meio Islands), with shrub and herbaceous vegetation (Deserta Island), with herbaceous vegetation (Siriú, Santana-de-Dentro and Santana-de-Fora Islands), and with shrub, herbaceous vegetation and cave (Batuta Island).

The presence of the otter on these islands can be related to favorable conditions for the animal in short periods of time throughout the year. It can be also related to the ability of the species to realize short distance movements from promontories to islands and from islands to islands, parallel to the coastline. Rivers, lakes and lagoons, located in mainland, might play an important role on these movements, as protection, fresh water and food availability to the animal. Although there are no fish inside the islands, neotropical otter should benefit from marine fishes along the coast of the islands. In this case the supralittoral zone can be used for resting between hunting activities.

Therefore, the presence of otters on these islands might be associated with the availability of supralittoral substrate, caves, freshwater and vegetation. Characteristics of the islands, combined with rivers, lakes and lagoons in mainland, might be representing a large ecological corridor for neotropical movements, facilitating the contact between subpopulations along the coast.

REFERENCES

- Alarcon, G.G., Simões-Lopes, P.C. (2004).** The neotropical otter *Lontra longicaudis* feeding habits in a marine coastal area, Southern Brazil. IUCN Otter Spec. Group Bull. **21(1)**: 24 - 30.
- Carvalho-Junior, O. (1990).** Aspectos da autoecologia de *Lutra longicaudis* (Olfers, 1818) no ecossistema da Lagoa do Peri, SC, Brasil. Tese de Dissertação em Hidroecologia. UFSC, Departamento de Biologia. 94 p.
- Carvalho-Junior, O., Banevicius, N.M.S., Mafra, E.O. (2004).** Distribution and characterization of environments used by otters in the coastal region of Santa Catarina State, Brazil. Journal of Coastal Research, 1087-1089.
- Carvalho-Junior, O. (2007).** No rastro da lontra Brasileira. Ed. Bernuncia, Florianópolis, Brasil, 112pp.
- Carvalho-Junior, O., Birolo, A.B., Macedo-Soares, L.C.P. de (2010a).** Ecological aspects of neotropical otter (*Lontra longicaudis*) in Peri Lagoon, South Brazil. IUCN Otter Spec. Group Bull, 27 (2): 105 - 115.
- Carvalho-Junior, O., Macedo-Soares, L.C.P. de, Birolo, A.B. (2010b).** Annual and interannual food habits variability of a neotropical otter (*Lontra longicaudis*) population in Conceição Lagoon, south of Brazil. IUCN Otter Spec. Group Bull. 27 (1): 24 - 32.
- Filippini, A. (2009).** Biogeografia dos Vertebrados de Ilhas de Santa Catarina: destaque em Aves Marinhas e Costeiras. Dissertação - Universidade Federal de Santa Catarina (UFSC). Florianópolis, SC, 351pp.
- Helder, J and Andrade, H.K. (1997).** Food and feeding habits of the neotropical river otter *Lontra*

longicaudis (Carnivora, Mustelidae). Mammalia, **61**: 193-203.

Pardini, R. (1998). Feeding ecology of the neotropical otter *Lontra longicaudis* in an Atlantic forest stream, south Brazil. Journal of Zoology, London, **245**: 385-391.

Quadros, J., Monteiro-Filho, E.L.A. (2001). Diet of the neotropical otter, *Lontra longicaudis*, in the Atlantic Forest, Santa Catarina State, Southern Brazil. Studies on neotropical fauna and environment. **36(1)**: 15-21.

Quintela, F.M., Porciúncula, R.A., Colares, E.P. (2008). Diet of *Lontra longicaudis* (Olfers) in a coastal stream in southern Rio Grande do Sul State, Brazil. Neotropical Biology and Conservation. **3(3)**: 119-125.

ACKNOWLEDGEMENTS - This work is sponsored by Petrobras, through Petrobras Environmental Program (Programa Petrobras Ambiental) and Conselho Federal dos Direitos Difusos (CFDD/MJ) - Fundo de Bens Lesados; and supported by Prefeitura Municipal de Florianópolis and IBAMA (Instituto Brasileiro de Meio Ambiente e dos Recursos Naturais Renováveis). We are most grateful to Gaëlle Moisan and Lachenal and Nathalie Eychenne, former ecovolunteers at the Otter Project, who kindly translate the abstract to French.

RÉSUMÉ

DISTRIBUTION DE LA LOUTRE A LONGUE QUEUE TITRE, *Lontra longicaudis* (OLFERS, 1818) (MUSTELIDAE), SUR LES ILES COTIERES DE SANTA CATARINA, SUD DU BRESIL

Ce travail présente la répartition de la Loutre à longue queue (*Lontra longicaudis*) sur les îles côtières de l'Etat de Santa Catarina. Vingt-trois îles ont été prospectées à partir de 46 circuits. La Loutre est présente sur 9 îles soit 39% des îles étudiées. Il est probable que *Lontra longicaudis* utilise ces îles comme étape durant ses déplacements d'un point à un autre. Ces résultats pourraient indiquer la présence d'un corridor écologique marin utilisé par la Loutre à longue queue.

RESUMEN

TITLE

Text

Resumo

DISTRIBUIÇÃO DA LONTRA NEOTROPICAL, *Lontra longicaudis* (OLFERS, 1818) (MUSTELIDAE) EM ILHAS COSTEIRAS DE SANTA CATARINA, SUL DO BRASIL

Este estudo apresenta a distribuição da lontra neotropical (*Lontra longicaudis*) em ilhas costeiras do Estado de Santa Catarina. O total de 23 ilhas foram amostradas em 46 expedições. Sinais de lontras foram encontrados em 9 ilhas, representando 39% do total. Acredita-se que a *Lontra longicaudis* utiliza estas ilhas como apoio para deslocamento de um lugar para o outro. Os resultados podem estar indicando a existência de um corredor ecológico utilizado pela lontra neotropical.